

# Bat Conservation Trust



Core Sustenance zones: Determining zone size reference list  
04.02.16

## *Myotis alcathoe*

Lucan, R., Andreas, M., Benda, P., Bartonicka, T., Brezinova, T., Hoffmannova, A., Hulova, S., Hulva, P., Neckarova, J., Reiter, A., Svacina, T., Salek, M., Horacek, I., 2009. .Alcathoe bat (*Myotis alcathoe*) in the Czech Republic: distributional status, roosting and feeding ecology. *Acta Chiropterologica* 11: 61-69.

## *Barbastella barbastellus*

Billington, G. 2000. Horner Woods Barbastelle Bat: radio tracking study. The National Trust.

Billington, G. 2012. Further research on the Barbastelle Bat Holnicote National Trust Estate, Exmoor, North Somerset. Greena Ecological Consultancy

Greenaway, F. 2008. Barbastelle Bats In The Sussex West Weald 1997 – 2008. Sussex Wildlife Trust/ West Weald Landscape Partnership

Greenaway, F., 2001. The barbastelle in Britain. *British Wildlife* 12: 327-334.

Hillen, J., Kiefer, A., Veith, M., 2009. Foraging site fidelity shapes the spatial organisation of a population of female western barbastelle bats. *Biological Conservation* 142: 817-823.

Hillen, J., Kiefer, A., Veith, M., 2010. Interannual fidelity to roosting habitat and flight paths by female western barbastelle bats. *Acta Chiropterologica* 12: 187-195.

Kerth, G., Melber, M., 2009. Species-specific barrier effects of a motorway on the habitat use of two threatened forest-living bat species. *Biological Conservation* 142: 270-279.

Murray, A., 2015. Unpublished data

Rush, T. & Billington, G. 2013. Report on a radio tracking study of Barbastelle bats at Hinkley Point C. Greena Ecological Consultancy.

Russo, D., Cistrone, L., Jones, G., Mazzoleni, S., 2004. Roost selection by barbastelle bats (*Barbastella barbastellus*, Chiroptera : Vespertilionidae) in beech woodlands of central Italy: consequences for conservation. *Biological Conservation* 117: 73-81.

Sierro, A., 1999. Habitat selection by barbastelle bats (*Barbastella barbastellus*) in the Swiss Alps (Valais). *Journal of Zoology* 248: 429-432.

Simon, M., Hüttenbügel, S., & Smit-Viergutz, J., 2004. Ecology and Conservation of Bats in Villages and Towns. Bundesamt für Naturschutz

# Bat Conservation Trust



- Steinhauser, D., Burger, F., Hoffmeister, U., Matez, G., Teige, T., Steinhauser, P., Wolz, I., 2002. Untersuchungen zur Ökologie der Mopsfledermaus, *Barbastella barbastellus* (Schreber, 1774), und der Bechsteinfledermaus, *Myotis bechsteinii* (Kuhl, 1817) im Süden des Landes Brandenburg. Schriftenr. Landschaftspflege. Naturschutz 71: 81–98.
- Zeale, M., Davidson-Watts, I., Jones, G., 2012. Home range use and habitat selection by barbastelle bats (*Barbastella barbastellus*): implications for conservation. Journal of Mammalogy 93: 1110-1118.

## *Myotis bechsteinii*

- Albrecht, K., Hammer, M. & Holzhaider, J. 2002: Telemetrische Untersuchungen zum Nahrungshabitatanspruch der Bechsteinfledermaus (*Myotis bechsteinii*) in Nadelwäldern bei Amberg in der Oberpfalz. — Schriftenreihe für Landschaftspflege und Naturschutz 71: 109–130.
- Bohnenstengel, T., 2012. Roost selection by the forest-dwelling bat *Myotis bechsteinii* (Mammalia: Chiroptera): implications for its conservation in managed woodlands. Bulletin de la Société Neuchâteloise des Sciences Naturelles. 132: 47-62.
- Dickins, E., Damant, C., 2015. Bernwood Forest, Buckinghamshire Summary and Review of Bechstein's Survey Results 2010-2014. Bernwood Environmental Conservation Services
- Dietz, M., Pir, J.B., 2009. Distribution and habitat selection of *Myotis bechsteinii* in Luxembourg: implications for forest management and conservation Folia Zoologica 58: 327-340.
- Fitzsimons, P., Hill, D., Greenaway, F., 2002. Bechstein's bat foraging habitat selection: Patterns of habitat use by female Bechstein's bats (*Myotis bechsteinii*) from a maternity colony in a British woodland. University of Bristol Report.
- Kerth, G., Melber, M., 2009. Species-specific barrier effects of a motorway on the habitat use of two threatened forest-living bat species. Biological Conservation 142: 270-279.
- Kerth, G., Morf, L., 2004. Behavioural and genetic data suggest that Bechstein's bats predominantly mate outside the breeding habitat. Ethology 110: 987-999.
- Kerth, G., Wagner, M., Konig, B., 2001. Roosting together, foraging apart: information transfer about food is unlikely to explain sociality in female Bechstein's bats (*Myotis bechsteinii*). Behavioural Ecology and Sociobiology 50: 283-291.
- Napal, M., Garin, I., Goiti, U., Salsamendi, E., Aihartza, J., 2013. Past deforestation of Mediterranean Europe explains the present distribution of the strict forest dweller *Myotis bechsteinii*. Forest Ecology and Management 293: 161-170.
- Palmer, E, 2014. Bechstein's and barbastelle bats in Gloucestershire and South Gloucestershire: a survey report for 2013-14

# Bat Conservation Trust



Palmer, E., Pimley, E., Sutton, G., Birks, J., 2013. A study on the population size, foraging range and roosting ecology of Bechstein's bats at Grafton Wood SSSI, Worcestershire. *Link Ecology and Swift Ecology*.

Schofield, H. & Morris, C. 2009. Report on the Bechstein's bat colony at Poplar Cottage, Balcombe, West Sussex. The Vincent Wildlife Trust.

Schofield, H., Morris, C., 2000. Ranging behaviour and habitat preferences of female Bechstein's bat, *Myotis bechsteinii* (Kuhl, 1818), in summer. The Vincent Wildlife Trust

Steinhauser, D., 2002. Untersuchungen zur Ökologie der Mopsfledermaus, *Barbastella barbastellus* (Schreber, 1774), und der Bechsteinfledermaus, *Myotis bechsteinii* (Kuhl, 1817), im Süden des Landes Brandenburg. In: Meschede, A., Heller, K.-G., Boye, P. (Eds.), *Ökologie, Wanderungen und Genetik von Fledermausen in Waldern*, vol. 71. Untersuchungen als Grundlage für den Fledermausschutz, Schriftenreihe für Landschaftspflege und Naturschutz, pp. 81–99.

## ***Plecotus auritus***

Entwistle, A., Racey, P., Speakman, J., 1996. Habitat exploitation by a gleaning bat, *Plecotus auritus*. *Philosophical Transactions of the Royal Society of London Series B – Biological Sciences* 351: 921-931.

Fuhrmann, M., Seitz, A., 1992. Nocturnal activity of the brown long-eared bat (*Plecotus auritus* L., 1758): data from radio-tracking in the Lenneberg forest near Maniz (Germany), in: I.G. Priede, S.M. Swift (Eds.), *Wildlife telemetry. Remote monitoring and tracking animals*. Ellis Horwood, New York, pp. 538-548.

Murphy, S., 2015. Unpublished data

Murphy, S., Greenaway, F., Hill, D., 2012. Patterns of habitat use by female brown long-eared bats presage negative impacts of woodland conservation management. *Journal of Zoology* 288: 177-183.

PLACE, 2002. Proceedings of the PLACE conference: Land Use, Ecology and Conservation of the Lower Derwent Valley.

Swift, S.M., Racey, P.A., 1983. Resource partitioning in two species of vespertilionid bats (Chiroptera) occupying the same roost. *Journal of Zoology* 200: 249-259.

## ***Myotis brandtii***

Berge, L., 2007. Resource partitioning between the cryptic species Brandt's bat (*Myotis brandtii*) and the whiskered bat (*M. mystacinus*) in the UK. Ph.D. thesis, University of Bristol.

# Bat Conservation Trust



## *Pipistrellus pipistrellus*

- Davidson-Watts, I., Jones, G., 2006. Differences in foraging behaviour between *Pipistrellus pipistrellus* (Schreber, 1774) and *Pipistrellus pygmaeus* (Leach, 1825). *Journal of Zoology* 268; 55-62.
- Eichstadt, H., Bassus, W., 1995. Untersuchungen zur Nahrungsökologie der Zwergfledermaus (*Pipistrellus pipistrellus*). *Nyctalus* 5: 561-584.
- Feyerabend, F., Simon, M., 2000. Use of roosts and roost switching in a summer colony of 45 kHz phonic type pipistrelle bats (*Pipistrellus pipistrellus* Schreber, 1774). *Myotis* 38: 51-59.
- Nicholls, B., Racey, P., 2006. Contrasting home-range size and spatial partitioning in cryptic and sympatric pipistrelle bats. *Behavioural Ecology and Sociobiology* 61: 131-142.

## *Myotis daubentonii*

- Dietz, M., Encarnacao, J., Kalko, E., 2006. Small scale distribution patterns of female and male Daubenton's bats (*Myotis daubentonii*). *Acta Chiropterologica* 8: 403-415.
- Encarnacao, J., 2012. Spatiotemporal pattern of local sexual segregation in a tree-dwelling temperate bat *Myotis daubentonii*. *Journal of Ethology* 30: 271-278.
- Kapfer, G., Rigot, T., Holsbeek, L. & Aron, S. (2008) Roost and hunting site fidelity of female and juvenile Daubenton's bat *Myotis daubentonii* (Kuhl, 1817) (Chiroptera: Vespertilionidae). *Mammalian Biology*, 73, 267–275.
- Lucan, R., Radil, J., 2010. Variability of foraging and roosting activities in adult females of Daubenton's bat (*Myotis daubentonii*) in different seasons. *Biologia* 65: 1072-1080.
- Parsons, K., Jones, G., 2003. Dispersion and habitat use by *Myotis daubentonii* and *Myotis nattereri* during the swarming season: implications for conservation. *Animal Conservation* 6: 283-290.
- Senior, P., Butlin, R.K., Altringham, J.D., 2005. Sex and segregation in temperate bats . *Proc. R. Soc. B*, 272: 2467–2473.

## *Rhinolophus ferrumequinum*

- Billington, G. 2000. Radio tracking study of greater horseshoe bats at Mells, Near Frome, Somerset. *English Nature*

# Bat Conservation Trust



- Billington, G. 2013. Cheddar Reservoir 2: Radio tracking studies of greater horseshoe and lesser horseshoe bats, June and August 2013. Greena Ecological Consultancy.
- Billington, G., 2002. Radio tracking study of Greater Horseshoe bats at Brockley Hall Stables Site of Special Scientific Interest. English Nature Research Report No. 442.
- Billington, G., 2003. Radio tracking study of greater horseshoe bats at Chudleigh Caves and Woods Site of Special Scientific Interest. English Nature Research Report No. 496
- Billington, G., 2004. Radio tracking study of greater horseshoe bats at Buckfastleigh Caves Site of Special Scientific Interest, 2003. English Nature Research Report No. 573.
- Bontadina, F. 2002. Conservation ecology in the horseshoe bats *Rhinolophus ferrumequinum* and *Rhinolophus hipposideros*. PhD Thesis, University of Bern.
- Collins, J., Billington, G., 2004. Monitoring (the Welsh colony of) Greater Horseshoe bats (*Rhinolophus ferrumequinum*) in the Wye Valley through radio-tracking and field survey to assess habitat radio-tracking and field survey to assess habitat use and condition. English Nature Research Report No. 495.
- Duverge, P., 1996. Foraging activity, habitat use, development of juveniles, and diet of the greater horseshoe bat (*Rhinolophus ferrumequinum* - Schreber 1774) in south-west England. PhD Thesis, University of Bristol.
- Flanders, J., Jones, G., 2009. Roost use, ranging behaviour and diet of greater horseshoe bats (*Rhinolophus ferrumequinum*) using a transitional roost. *Journal of Mammalogy* 90: 888-896.
- Jones, G. and Morton, M., 1992. Radio-tracking studies on habitat use by greater horseshoe bats (*Rhinolophus ferrumequinum*) pp 521–537 in Wildlife telemetry: Remote monitoring and tracking of animals. Priede, I. G. and S. M. Swift. Ellis Horwood. Chichester, United Kingdom.
- Jones, G., Billington, G., 1999. Radio tracking study of greater horseshoe bats at Cheddar, North Somerset. English Nature.
- Jones, G., Duverge, P., Ransome, R., 1995. Conservation biology of an endangered species: Field studies of greater horseshoe bats. *Journal of the Zoological Society of London*, 67: 309-324.
- Robinson, M., Webber, M., Stebbings, R., 2000. Dispersal and foraging behaviour of greater horseshoe bats, Brixham, Devon. English Nature Research Report No. 344.
- Rossiter, S.J., Jones, G., Ransome, R.D., Barratt, E.M., 2002. Relatedness structure and kin-biased foraging in the greater horseshoe bat (*Rhinolophus ferrumequinum*). *Behavioural Ecology and Sociobiology* 51: 510-518.

# Bat Conservation Trust



## *Plecotus austriacus*

Razgour O., Whitby D., Dahlberg E., Barlow K., Hanmer J., Haysom K., McFarlane H., Wicks L., Williams C., Jones G., 2013. Conserving grey long-eared bats (*Plecotus austriacus*) in our landscape: a conservation management plan. Available to download from the Bat Conservation Trust website [www.bats.org.uk](http://www.bats.org.uk)

Razgour, O., Hanmer, J., Jones, G., 2011. Using multi-scale modelling to predict habitat suitability for species of conservation concern: The grey long-eared bat as a case study. *Biological Conservation* 144: 2922-2930.

## *Nyctalus leisleri*

Haddow, J. 2014. Unpublished data

Shiel, C., Shiel, R., Fairley, J., 1999. Seasonal changes in the foraging behaviour of Leisler's bats (*Nyctalus leisleri*) in Ireland as revealed by radio-telemetry. *Journal of Zoology* 249: 347-358.

Szentkuti, S., Bontadina, F., Spada, M., Moretti, M., Zambelli, N., Martinoli, A., Arlettaz, R., 2013. Factors underlying migratory bat aggregations in chestnut groves. *Endangered Species Research* 21: 105-114.

Waters, D., Jones, G., Furlong, M., 1999. Foraging ecology of Leisler's bat (*Nyctalus leisleri*) at two sites in southern Britain. *Journal of Zoology* 249: 173-180.

## *Rhinolophus hipposideros*

Billington, G. 2005. Radio tracking study of lesser horseshoe bats at Hestercombe House Site of Special Scientific Interest, July 2005. English Nature Somerset & Gloucestershire Team.

Billington, G. 2013. Cheddar Reservoir 2: Radio tracking studies of greater horseshoe and lesser horseshoe bats, June and August 2013. Greena Ecological Consultancy.

Billington, G., 2001. Report on a radio tracking study of lesser horseshoe bats associated with the A487 Llanwnda to South of Llanllfni Road Improvement. Greena Ecological Consultancy.

Bontadina, F., Schofield, H., Naef-Daenzer, B., 2002. Radio-tracking reveals that lesser horseshoe bats (*Rhinolophus hipposideros*) forage in woodland. *Journal of Zoology* 258: 281-290.

# Bat Conservation Trust



- Duvergé, L. 2008. Report on bat surveys carried out at Hestercombe House SSSI Taunton, Somerset, in 2007 and 2008. Kestrel Wildlife Consultants.
- Holzhaider, J., Kriner, E., Rudolph, B.-U., Zahn, A., 2002. Radio-tracking a lesser horseshoe bat (*Rhinolophus hipposideros*) in Bavaria: an experiment to locate roosts and foraging sites. *Myotis* 40: 47-54.
- Knight, T. 2006. The use of landscape features and habitats by the lesser horseshoe bat (*Rhinolophus hipposideros*). PhD Thesis, University of Bristol.
- Knight, T., Jones, G., 2009. Importance of night roosts for bat conservation: roosting behaviour of the lesser horseshoe bat *Rhinolophus hipposideros*. *Endangered Species Research* 9: 79-86.
- Schofield, H., Morris, C., 2000. Ranging behaviour and habitat preferences of female Bechstein's bat, *Myotis bechsteinii* (Kuhl, 1818), in summer. The Vincent Wildlife Trust.
- Zahn, A., Holzhaider, J., Kriner, E., Maier, A., Kayikcioglu, A., 2008. Foraging activity of *Rhinolophus hipposideros* on the island of Herrenchiemsee, Upper Bavaria. *Mammalian Biology* 73: 222-229

## *Pipistrellus nathusii*

- Flaquer, C., Puig-Montserrat, X., Goiti, U., Vidal, F., Curcó, A., Russo, D., 2009. Habitat Selection in Nathusius' Pipistrelle (*Pipistrellus nathusii*): The Importance of Wetlands. *Acta Chiropterologica* 11: 149-155.
- Hargreaves, D., 2011. Nathusius' pipistrelle tracking report and update map.

## *Myotis nattereri*

- Lundy, M., Buckley, D., Boston, E., et al. 2012. Behavioural context of multi-scale species distribution models assessed by radio-tracking. *Basic and Applied Ecology* 13: 188-195.
- Meschede, A., Heller, K.G., 2000. Ökologie und Schutz von Fledermäusen in Wäldern. Landwirtschaftsverlag, Münster.
- Parsons, K., Jones, G., 2003. Dispersion and habitat use by *Myotis daubentonii* and *Myotis nattereri* during the swarming season: implications for conservation. *Animal Conservation* 6: 283-290.
- Siemers, B., Kaipf, I., Schnitzler, H., 1999. The use of day roosts and foraging grounds by Natterer's bats (*Myotis nattereri* Kuhl, 1818) from a colony in southern Germany. *Zeitschrift Fur Saugetierkunde*, 64: 241–245 .
- Smith, P. 2001. Habitat preferences, range use and roosting ecology of Natterer's bat *Myotis nattereri* in a grassland-woodland landscape. Ph.D. thesis, University of Aberdeen.

# Bat Conservation Trust



- Smith, P., Racey, P., 2005. The itinerant Natterer: physical and thermal characteristics of summer roosts of *Myotis nattereri* (Mammalia : Chiroptera). Journal of Zoology 266: 171-180.
- Smith, P., Racey, P., 2008. Natterer's bats prefer foraging in broad-leaved woodlands and river corridors. Journal of Zoology 275: 314-322.
- Trappmann, C., Clemen, G., 2001. Observations on exploitation of the hunting territory of Natterer's bat *Myotis nattereri* (Kuhl, 1817) by means of telemetry. Acta Biologica Benrodis 11: 1-31.
- Zeale et al. 2014. Improving mitigation success where bats occupy houses and historic buildings, particularly churches. Defra Research Project WM0322.

## *Nyctalus noctula*

- Kronwitter, F., 1988. Population structure, habitat use and activity patterns of the noctule bat *Nyctalus noctula* Schreb., 1774 (Chiroptera: Vespertilionidae) revealed by radio-tracking. Myotis 26: 23-85.
- Mackie, I., Racey, P., 2007. Habitat use varies with reproductive state in noctule bats (*Nyctalus noctula*): Implications for conservation. Biological Conservation 140: 70-77.
- Meschede, A., Heller, K.G., 2000. Ökologie und Schutz von Fledermäusen in Wäldern. Landwirtschaftsverlag, Münster.
- Schmidt, A., 1988. Beobachtungen zur Lebensweise des Abendseglers, *Nyctalus noctula* (Schreber, 1774), im Süden des Bezirkes Frankfurt/O. Nyctalus 5: 389-422.

## *Eptesicus serotinus*

- Catto, C.M.C., Hutson, A.M., Racey, P.A., Stephenson, P.J., 1996. Foraging behaviour and habitat use of the serotine bat (*Eptesicus serotinus*) in southern England. Journal of Zoology, 238: 623-633.
- Cornes, B., 2014. Unpublished data.
- Dyer, S., 2014. Unpublished data.
- Harbusch, C., 2003. Aspects of the ecology of serotine bats (*Eptesicus serotinus*, Schreber 1774) in contrasting landscapes in Southwest Germany and Luxembourg. Ph.D. thesis, University of Aberdeen.
- Hargeaves, D., 2011. Serotine bat radio tracking report.

# Bat Conservation Trust



Kervyn, T., Brasseur, J., Libois, R., 1997. Utilisation de l'habitat par la sérotine commune *Eptesicus serotinus* en Lorraine Belge. Bulletin de la Société Neuchâteloise des Sciences Naturelles, 120: 35-41.

Robinson, M., Stebbings, R., 1997. Home range and habitat use by the serotine bat, *Eptesicus serotinus*, in England. Journal of Zoology 243: 117-136.

Rosenau, S., 2001. Untersuchungen zur Quartiernutzung und Habitatnutzung der Breitflügelfledermaus *Eptesicus serotinus* (Schreber, 1774) im Berliner Stadtgebiet (Bezirk Spandau). FU Berlin.

Simon, M., Hüttenbügel, S., Smit-Viergutz, J., 2004. Ökologie und Schutz von Fledermäusen in Dörfern und Städten, In Schriftenreihe für Landschaftspflege und Naturschutz 76. Bundesamtes für Naturschutz, Münster.

## *Pipistrellus pygmaeus*

Bartonicka, T., Bielik, A., Rehak, Z., 2008. Roost switching and activity patterns in the soprano pipistrelle, *Pipistrellus pygmaeus*, during lactation. Annales Zoologici Fennici, 45: 503-512.

Davidson-Watts, I., Jones, G., 2006. Differences in foraging behaviour between *Pipistrellus pipistrellus* (Schreber, 1774) and *Pipistrellus pygmaeus* (Leach, 1825). Journal of Zoology 268: 55-62.

Nicholls, B., Racey, P., 2006. Contrasting home-range size and spatial partitioning in cryptic and sympatric pipistrelle bats. Behavioural Ecology and Sociobiology 61: 131-142.

Ryan, M. 2015. Unpublished data

## *Myotis mystacinus*

Berge, L., 2007. Resource partitioning between the cryptic species Brandt's bat (*Myotis brandtii*) and the whiskered bat (*M. mystacinus*) in the UK. Ph.D. thesis, University of Bristol.

Buckley, D., Lundy, M., Boston, E., Scott, D., Gager, Y., Prodöhl, P., Marnell, F., Montgomery W.I., Teeling, E., 2013. The spatial ecology of the whiskered bat (*Myotis mystacinus*) at the western extreme of its range provides evidence of regional adaptation. Mammalian Biology 78: 198-204.